(19) World Intellectual Property Organization International Bureau



. 1948 WINDS 17 EIRIN BIND 1911 (1) IN SIN CITI SWINDING CON BIND CON CINING CONTING CON CINING CON

(43) International Publication Date 5 July 2001 (05.07.2001)

PCT

(10) International Publication Number WO 01/47509 A2

(51) International Patent Classification7:

.____

A61K 31/00

- (21) International Application Number: PCT/GB00/05006

(22) International Filing Date: 22 December 2000 (22.12.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

9930688.8 24 December 1999 (24.12.1999) GB 9930689.6 24 December 1999 (24.12.1999) GB 9930690.4 24 December 1999 (24.12.1999) GB 9930692.0 24 December 1999 (24.12.1999) GB

- (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): BUCKINGHAM, Robin, Edward [GB/GB]; SmithKline Beecham Pahrmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).

- (74) Agent: RUTTER, Keith; Corporate Intellectual Property, SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

 Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

509 A

(54) Title: NOVEL METHOD OF TREATMENT

(57) Abstract: A method for the treatment of diabetes mellitus, especially Type II diabetes and the cardiac conditions associated with diabetes mellitus in a mammal such as a human, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus.

10

15

20

25

30

35

40

NOVEL METHOD OF TREATMENT

This invention relates to a method of treatment, in particular to a method for the treatment of diabetes mellitus, especially non-insulin dependent diabetes (NIDDM) or Type II diabetes, and the cardiac conditions associated with diabetes mellitus.

It is known that a common sequela of the Type II diabetes syndrome is the development of cardiac conditions. In particular, it is known that Type II diabetes can lead to the development of heart failure, including congestive heart failure (also known as chronic heart failure).

Agents used in the treatment of heart failure, including congestive heart failure, include endothelin antagonists, beta-blockers, ACE inhibitors and diuretics.

Endothelin is a potent vasoconstrictor peptide synthesized and released by the vascular endothelium. International Patent Applications Publication Numbers WO 93/08799 and WO 94/25013 (SmithKline Beecham Corporation) disclose certain indanes and indenes as endothelin receptor antagonists. International Patent Application Publication Number WO 97/04772 (SmithKline Beecham Corporation) discloses certain pyrroles, pyrazoles, and triazoles as endothelin receptor antagonists.

Beta-blockers are compounds that act as competitive antagonists at beta-adrenergic receptor sites and include the compounds disclosed in reference texts such as Martindale 32nd Edition "The Complete Drug Reference" especially page 777. The alkanolamines are well known examples of beta-blockers. The alkanolamines are used in the treatment of cardiovascular disorders such as hypertension, angina pectoris, cardiac arrhythmias, and myocardial infarction.

Angiotensin converting enzyme (ACE) inhibitors are compounds that are used in the treatment of heart failure, hypertension, and myocardial infarction. And include the compounds disclosed in reference texts such as Martindale 32nd Edition "The Complete Drug Reference", especially page 776.

Diuretics are compounds which promote the excretion of urine and thus a reduction in blood plasma volume and include those compounds disclosed in reference texts such as Martindale 32nd Edition The Complete Drug Reference, especially page 778. Diuretic compounds may be divided into classes according to their mode of action. These classes include carbonic anhydrase inhibitors, loop diuretics, potassium-sparing diuretics, and thiazides

European Patent Application, Publication Number 0,306,228 relates to certain thiazolidinedione derivatives disclosed as having antihyperglycaemic and antihyperlipidaemic activity. One particular thiazolidinedione disclosed in EP 0306228 is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-

WO 01/47509 PCT/GB00/05006

2,4-dione (hereinafter 'Compound (I)'). WO 94/05659 discloses certain salts of Compound (I) including the maleate salt at example 1 thereof.

Compound (I) is an example of a class of anti-hyperglycaemic agents known as 'insulin sensitisers'. In particular Compound (I) is a thiazolidinedione insulin sensitiser.

European Patent Applications, Publication Numbers: 0008203, 0139421, 0032128, 0428312, 0489663, 0155845, 0257781, 0208420, 0177353, 0319189, 0332331, 0332332, 0528734, 0508740; International Patent Application, Publication Numbers 92/18501, 93/02079, 93/22445 and United States Patent Numbers 5104888 and 5478852, also disclose certain thiazolidinedione insulin sensitisers.

Another series of compounds generally recognised as having insulin sensitiser activity are those typified by the compounds disclosed in International Patent Applications, Publication Numbers WO 93/21166 and WO 94/01420. These compounds are herein referred to as 'acyclic insulin sensitisers'. Other examples of acyclic insulin sensitisers are those disclosed in United States Patent Number 5232945 and International Patent Applications, Publication Numbers WO 92/03425 and WO 91/19702. Further examples of insulin sensitiser are those disclosed in WO 97/31907 and GW262570.

Examples of other insulin sensitisers are those disclosed in European Patent Application, Publication Number 0533933, Japanese Patent Application Publication Number 05271204 and United States Patent Number 5264451.

The above mentioned publications are incorporated herein by reference.

It is now considered that Compound (I) in combination with an agent used in the treatment of heart failure, including congestive heart failure, provides a beneficial effect upon glycaemic control and ameliorates the cardiac conditions associated with diabetes mellitus, especially that associated with the onset and development of heart failure, beyond a mere additive effect. Such a combination is therefore particularly useful for the treatment of diabetes mellitus, especially Type II diabetes and the cardiac conditions arising from diabetes mellitus. The treatment is also indicated to proceed with minimum side effects.

Accordingly, the invention provides a method for the treatment of diabetes mellitus, especially Type II diabetes and the cardiac conditions associated with diabetes mellitus in a mammal such as a human, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus.

Cardiac conditions associated with diabetes mellitus includes those cardiac conditions arising from diabetes mellitus.

5

10

15

20

25

30

10

15

20

25

30

Cardiac conditions arising from diabetes mellitus include heart failure, for example congestive heart failure.

A suitable agent used in the treatment of cardiac conditions associated with diabetes mellitus, such as heart failure, includes a beta-blocker, an ACE inhibitor or a diuretic. A suitable agent used in the treatment of cardiac conditions associated with diabetes mellitus, such as heart failure, also includes an endothelin antagonist.

In another aspect the invention provides an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus, for use in a method for the treatment of diabetes mellitus, especially Type II diabetes, and the cardiac conditions associated with diabetes mellitus.

The method comprises either co-administration of an insulin sensitiser, such as Compound (I), and the agent used in the treatment of the cardiac conditions associated with diabetes mellitus or the sequential administration thereof.

Co-administration includes administration of a formulation which includes both an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus or the essentially simultaneous administration of separate formulations of each agent.

In another aspect the invention provides the use of an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus, for use in the manufacture of a composition for the treatment of diabetes mellitus, especially Type II diabetes and the cardiac conditions associated with diabetes mellitus.

A suitable thiazolidinedione insulin sensitiser is Compound (I). Other suitable thiazolidinedione insulin sensitisers include (+)-5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl]thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl]thiazolidine-2,4-dione (or pioglitazone), and 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl)thiazolidine-2,4-dione (or englitazone).

Suitable endothelin antagonists include those disclosed in WO 93/08799,
WO 94/25013, and WO 97/04772.
Suitable beta-blockers include acebutolol, alprenolol, amosulatol, arotinolol, atenolol, befunolol, betaxolol, bevantolol, bisoprolol, bopindolol, bucindolol, bufetolol, bufuralol, bunitrolol, bupranolol, carazolol, carteolol, carvedilol, celiprolol, chloranolol, dilevalol, epanolol, esmolol, flestolol, indenolol, labetalol, levobunolol, levomoprolol, medroxalol, mepindolol, metipranolol, metoprolol,

10

15

20

25

30

35

40

nadolol, nebivolol, nipradilol, oxprenolol, penbutolol, pindolol, practolol, propranolol, sotalol, talinolol, tertatolol, tilisolol, and timolol.

Suitable ACE inhibitors include alacepril, benazepril, captopril, ceronapril, cilazepril, delapril, enalapril, enalaprilat, fosinopril, imidapril, libenzapril, lisinopril, moexipril, moveltipril, pentopril, perindopril, quinapril, ramipril, spirapril, temocapril, teprotide, trandolapril, and zofenopril.

Suitable diuretics include acetazolamide, brinzolamide, dichlorphenamide, dorzolamide, methazolamide, azosemide, bumetanide, ethacrynic acid, etozolin, frusemide, piretanide, torasemide, isosorbide, mannitol, amiloride, canrenoate potassium, canrenone, spironolactone, triamterene, althiazide, bemetizide, bendrofluazide, benzthiazide, buthiazide, chlorothiazide, chlorothiazide, chlorothiazide, cyclopenthiazide, cyclothiazide, epithiazide, hydrochlorothiazide, hydroflumethiazide, indapamide, mebutizide, mefruside, methylclothiazide, meticrane, metolazone, polythiazide, quinethazone, teclothiazide, trichlormethiazide, tripamide, and xipamide.

In one particular aspect, the method comprises the administration of up to 12mg such as 2 to 12 mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 2 to 4, 4 to 8 or 8 to 12 mg of Compound (I) per day.

Particularly, the method comprises the administration of 2 to 4mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 4 to 8mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 8 to 12 mg of Compound (I), especially when administered per day.

Preferably, the method comprises the administration of 2 mg of Compound (I), especially when administered per day.

Preferably, the method comprises the administration of 4 mg of Compound (I), especially when administered per day.

Preferably, the method comprises the administration of 8 mg of Compound (I), especially when administered per day.

It will be understood that the insulin sensitiser, such as Compound (I) and the agent used in the treatment of the cardiac conditions associated with diabetes mellitus such as the endothelin antagonist, the beta-blocker, the ACE inhibitor, or the diuretic are each administered in a pharmaceutically acceptable form, including pharmaceutically acceptable derivatives such as pharmaceutically acceptable salts, esters and solvates thereof, as appropriate of the relevant pharmaceutically active agent. In certain instances herein the names used for the relevant agent used in the treatment of the cardiac conditions associated with diabetes mellitus may relate to a particular pharmaceutical form of the relevant

10

15

20

25

30

35

40

active agent. It will be understood that all pharmaceutically acceptable forms of the active agents per se are encompassed by this invention.

Suitable pharmaceutically acceptable salted forms of Compound (I) include those described in EP 0306228 and WO94/05659. A preferred pharmaceutically acceptable salt is a maleate.

Suitable pharmaceutically acceptable solvated forms of Compound (I) include those described in EP 0306228 and WO94/05659, in particular hydrates.

Suitable pharmaceutically acceptable forms of other active agents including the agent used in the treatment of the cardiac conditions associated with diabetes mellitus depend upon the particular agent used but include known pharmaceutically acceptable forms of the particular agent chosen, for example those disclosed in the above mentioned publications.

Compound (I) or, a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, may be prepared using known methods, for example those disclosed in EP 0306228 and WO94/05659. The disclosures of EP 0306228 and WO94/05659 are incorporated herein by reference.

Certain of the active agents including the thiazolidinedione insulin sensitisers such as Compound (I) may exist in one of several tautomeric forms, all of which are encompassed in this invention as individual tautomeric forms or as mixtures thereof. Certain of the active agents mentioned herein, including Compound (I) contain one or more chiral carbon atom, and hence can exist in distinct stereoisomeric forms, the present invention encompasses all of these isomeric forms whether as individual isomers or as mixtures of isomers, including racemates.

The particular method of preparation of the active agent used in the invention will depend upon the agent chosen but will in general be selected from methods known in the art.

The endothelin antagonist of choice is prepared according to known methods, for example those disclosed in WO 93/08799, WO 94/25013, and WO 97/04772.

The beta-blocker of choice is prepared according to known methods, such methods are found or are referred to in standard reference texts, such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale 32nd Edition The Complete Drug Reference (London, The Pharmaceutical Press) or the above mentioned publications.

The ACE inhibitor of choice is prepared according to known methods, such methods are found or are referred to in standard reference texts, such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale 32nd Edition The Complete Drug Reference (London, The Pharmaceutical Press) or the above mentioned publications.

WO 01/47509 PCT/GB00/05006

The diuretic of choice is prepared according to known methods, such methods are found or are referred to in standard reference texts, such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale 32nd Edition The Complete Drug Reference (London, The Pharmaceutical Press) or the above mentioned publications.

As used herein the term 'pharmaceutically acceptable' embraces both human and veterinary use: for example the term 'pharmaceutically acceptable' embraces a veterinarily acceptable compound.

For the avoidance of doubt, when reference is made herein to scalar amounts, including mg amounts, of Compound (I) in a pharmaceutically acceptable form, the scalar amount referred to is made in respect of Compound (I) per se. For example 2 mg of Compound (I) in the form of the maleate salt is that amount of maleate salt which contains 2 mg of Compound (I).

Diabetes mellitus is preferably Type II diabetes.

The term 'cardiac conditions' includes heart failure.

Glycaemic control may be characterised using conventional methods, for example by measurement of a typically used index of glycaemic control such as fasting plasma glucose or glycosylated haemoglobin (Hb A1c). Such indices are determined using standard methodology, for example those described in: Tuescher A, Richterich, P., Schweiz. med. Wschr. 101 (1971), 345 and 390 and Frank P., 'Monitoring the Diabetic Patent with Glycosolated Hemoglobin Measurements', Clinical Products 1988.

In the method of the invention, the active medicaments are preferably administered in pharmaceutical composition form. As indicated above, such compositions can include both medicaments or one only of the medicaments. Accordingly, in one aspect the present invention also provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) especially 2 to 12 mg thereof, an agent used in the treatment of the cardiac conditions associated with diabetes mellitus and a pharmaceutically acceptable carrier therefor.

Such compositions may be prepared by admixing an insulin sensitiser, such as Compound (I) especially 2 to 12 mg thereof, the agent used in the treatment of the cardiac conditions associated with diabetes mellitus and a pharmaceutically acceptable carrier therefor.

Usually the compositions are adapted for oral administration. However, they may be adapted for other modes of administration, for example parenteral administration, sublingual or transdermal administration.

The compositions may be in the form of tablets, capsules, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid preparations, such as oral or sterile parenteral solutions or suspensions.

5

10

15

20

25

30

35

10

15

20

In order to obtain consistency of administration it is preferred that a composition of the invention is in the form of a unit dose.

Unit dose presentation forms for oral administration may be tablets and capsules and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize starch, calcium phosphate, sorbitol or glycine; tabletting lubricants, for example magnesium stearate; disintegrants, for example starch, polyvinylpyrrolidone, sodium starch glycollate or microcrystalline cellulose; or pharmaceutically acceptable wetting agents such as sodium lauryl sulphate.

The compositions are preferably in a unit dosage form in an amount appropriate for the relevant daily dosage.

Suitable dosages including unit dosages of the Compound of formula (I) comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 mg of Compound (I).

In the treatment the medicaments may be administered from 1 to 6 times a day, but most preferably 1 or 2 times per day.

Particular dosages of Compound (I) are 2mg/day, 4mg/day, including 2mg twice per day, and 8 mg/day, including 4mg twice per day.

Suitable dosages of the agent used in the treatment of the cardiac conditions associated with diabetes mellitus are those used in the art for the particular agent chosen.

Suitable dosages of the endothelin antagonist depend on the endothelin antagonist chosen, but include those disclosed in WO 93/08799, WO 94/25013, and WO 97/04772.

Suitable dosages of the beta-blocker, such as the alkanolamine, include the known dosages, including unit doses, for these compounds as described or referred to in reference text such such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale 32nd Edition The Complete Drug Reference (London, The Pharmaceutical Press), or the above mentioned publications.

Suitable dosages of the ACE inhibitor include the known dosages, including unit doses, for these compounds as described or referred to in reference text such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale 32nd Edition The Complete Drug Reference (London, The Pharmaceutical Press) or the charmaceutical press.

Reference (London, The Pharmaceutical Press) or the above mentioned publications.

Suitable dosages including unit dosages of the diuretic include the known dosages including unit doses for these compounds as described or referred to in reference text such as the British and US Pharmacopoeias, Remington's

WQ 01/47509 PCT/GB00/05006

Pharmaceutical Sciences (Mack Publishing Co.), Martindale 32nd Edition The Complete Drug Reference (London, The Pharmaceutical Press).

The solid oral compositions may be prepared by conventional methods of blending, filling or tabletting. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are of course conventional in the art. The tablets may be coated according to methods well known in normal pharmaceutical practice, in particular with an enteric coating.

Oral liquid preparations may be in the form of, for example, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel, hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid; and if desired conventional flavouring or colouring agents.

For parenteral administration, fluid unit dosage forms are prepared utilizing the compound and a sterile vehicle, and, depending on the concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, a preservative and buffering agent can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the Compound (I) suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

Compositions may contain from 0.1% to 99% by weight, preferably from 10-60% by weight, of the active material, depending upon the method of administration.

Composition may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

The compositions are prepared and formulated according to conventional methods, such as those disclosed in standard reference texts, for example the

5

10

15

20

25

30

35

15

20

British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Complete Drug Reference 32nd Edition and Harry's Cosmeticology (Leonard Hill Books) or the above mentioned publications.

The present invention also provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) especially 2 to 12 mg thereof, an agent used in the treatment of the cardiac conditions associated with diabetes mellitus and a pharmaceutically acceptable carrier therefor, for use as an active therapeutic substance.

In particular, the present invention provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) especially 2 to 12 mg thereof, an agent used in the treatment of the cardiac conditions associated with diabetes mellitus and a pharmaceutically acceptable carrier therefor, for use in the treatment of diabetes mellitus, especially Type II diabetes and the cardiovascular conditions arising from diabetes mellitus.

A range of 2 to 4mg includes a range of 2.1 to 4, 2.2 to 4, 2.3 to 4, 2.4 to 4, 2.5 to 4, 2.6 to 4, 2.7 to 4, 2.8 to 4, 2.9 to 4 or 3 to 4mg.

A range of 4 to 8mg includes a range of 4.1 to 8, 4.2 to 8, 4.3 to 8, 4.4 to 8, 4.5 to 8, 4.6 to 8, 4.7 to 8, 4.8 to 8, 4.9 to 8, 5 to 8, 6 to 8 or 7 to 8mg.

A range of 8 to 12 mg includes a range of 8.1 to 12, 8.2 to 12, 8.3 to 12, 8.4 to 12, 8.5 to 12, 8.6 to 12, 8.7 to 12, 8.8 to 12, 8.9 to 12, 9 to 12, 10 to 12 or 11 to 12mg.

No adverse toxicological effects have been established for the compositions or methods of the invention in the above mentioned dosage ranges.

Claims:

5

15

25

30

- 1. A method for the treatment of diabetes mellitus, especially Type II diabetes and the cardiac conditions associated with diabetes mellitus in a mammal such as a human, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus.
- 2. A method according to claim 1, wherein the agent used in the treatment of cardiac conditions associated with diabetes mellitus is a beta-blocker, an ACE inhibitor or a diuretic.
 - 3. A method according to claim 1, wherein the agent used in the treatment of cardiac conditions associated with diabetes mellitus is an endothelin antagonist.
 - 4. A method according to any one of claims 1 to 3, wherein the insulin sensitiser is s thiazolidinedione insulin sensitiser.
- 5. A method according to any one of claims 1 to 4, wherein the thiazolidinedione insulin sensitiser is Compound (I).
 - 6. A method according to any one of claims 1 to 4, wherein the thiazolidinedione insulin sensitiser is selected from the list consisting of: (+)-5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl]thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl]thiazolidine-2,4-dione (or pioglitazone), and 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl)thiazolidine-2,4-dione (or englitazone).
 - 7. A method according to claim 1 or claim 2, wherein the beta-blocker is selected from the list consisting of acebutolol, alprenolol, amosulatol, arotinolol, atenolol, befunolol, betaxolol, bevantolol, bisoprolol, bopindolol, bucindolol, bufetolol, bufuralol, bunitrolol, bupranolol, carazolol, carteolol, carvedilol, celiprolol, chloranolol, dilevalol, epanolol, esmolol, flestolol, indenolol, labetalol, levobunolol, levomoprolol, medroxalol, mepindolol, metipranolol, metoprolol, nadolol, nebivolol, nipradilol, oxprenolol, penbutolol, pindolol, practolol, propranolol, sotalol, talinolol, tertatolol, tilisolol and timolol.

10

- 8. A method according to claim 1 or claim 2, wherein the ACE inhibitor blocker is selected from the list consisting of alacepril, benazepril, captopril, ceronapril, cilazepril, delapril, enalapril, enalaprilat, fosinopril, imidapril, libenzapril, lisinopril, moexipril, moveltipril, pentopril, perindopril, quinapril, ramipril, spirapril, temocapril, teprotide, trandolapril and zofenopril.
- 9. A method according to claim 1 or claim 2, wherein the diuretics is selected from the list consisting of acetazolamide, brinzolamide, dichlorphenamide, dorzolamide, methazolamide, azosemide, bumetanide, ethacrynic acid, etozolin, frusemide, piretanide, torasemide, isosorbide, mannitol, amiloride, canrenoate potassium, canrenone, spironolactone, triamterene, althiazide, bemetizide, bendrofluazide, benzthiazide, buthiazide, chlorothiazide, chlorthalidone, clopamide, cyclopenthiazide, cyclothiazide, epithiazide, hydrochlorothiazide, hydroflumethiazide, indapamide, mebutizide, mefruside, methylclothiazide, meticrane, metolazone, polythiazide, quinethazone, teclothiazide, trichlormethiazide, tripamide and xipamide.
- 10. A method according to any one of claims 1 to 9 which comprises either co-administration of an insulin sensitiser and the agent used in the treatment of the cardiac conditions associated with diabetes mellitus or the sequential administration thereof.

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 5 July 2001 (05.07.2001)

PCT

(10) International Publication Number WO 01/047509 A3

(51) International Patent Classification?: A61K 45/06, A61P 3/10

(21) International Application Number: PCT/GB00/05006

(22) International Filing Date:

22 December 2000 (22.12.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

 9930688.8
 24 December 1999 (24.12.1999)
 GB

 9930689.6
 24 December 1999 (24.12.1999)
 GB

 9930690.4
 24 December 1999 (24.12.1999)
 GB

 9930692.0
 24 December 1999 (24.12.1999)
 GB

- (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): BUCKINGHAM, Robin, Edward [GB/GB]; GlaxoSmithKline Pahrmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).

- (74) Agent: RUTTER, Keith; GlaxoSmithKline, Corporate Intellectual Property (CN9.25.1), 980 Great West Road, Brentford, Middlesex TW8 9GS (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- (88) Date of publication of the international search report: 12 September 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITION FOR THE TREATMENT DIABETES MELLITUS CONTAINING AN INSULINE SENSITIZER AND AGENT USED IN THE TREATMENT OF CARDIAC CONDITIONS

(57) Abstract: A method for the treatment of diabetes mellitus, especially Type II diabetes and the cardiac conditions associated with diabetes mellitus in a mammal such as a human, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, such as Compound (I), and an agent used in the treatment of the cardiac conditions associated with diabetes mellitus.

01/047509 A3

PCT/GB 00/05006

CLASSIFICATION OF SUBJECT MATTER
C 7 A61K45/06 A61F A61P3/10 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED finimum documentation searched (classification system followed by classification symbols) IPC 7 A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, EPO-Internal, CHEM ABS Data, CANCERLIT, MEDLINE, EMBASE, WPI Data C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. X SHIMOYAMA, MASAKI ET AL: "Hemodynamic 1,2,4-7, basis for the acute cardiac effects of 10 troglitazone in isolated perfused rat hearts. DIABETES, (MARCH, 1999) VOL. 48, NO. 3, PP. 609-615. XP001035237 page 610, column 2, paragraph 5 -page 611, paragraph 2 X,P WO 00 27434 A (ARCH JONATHAN ROBERT 1,2,4-7, SANDERS ; SMITHKLINE BEECHAM PLC (GB)) 18 May 2000 (2000-05-18) claims -/--Further documents are listed in the continuation of box C. X Patent family members are listed in annex. Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relevance; the claimed Invention cannot be considered to involve an inventive step when the document is combined with one or more other such docu- document referring to an oral disclosure, use, exhibition or other means ments, such combination being obvious to a person skilled document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the International search Date of mailing of the international search report 2 **%** < 04 2002 11 April 2002 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016 Leherte, C

Form PCT/ISA/210 (second sheet) (July 1992)

PCT/GB 00/05006

	tion) DOCUMENTS CONSIDERED TO BE RELEVANT	Polomet to aloim No
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DAGOGO-JACK S ET AL: "PATHOPHYSIOLOGY OF TYPE 2 DIABETES AND MODES OF ACTION OF THERAPEUTIC INTERVENTIONS" ARCHIVES OF INTERNAL MEDICINE, XX, XX, vol. 157, no. 16, 8 September 1997 (1997-09-08), pages 1802-1817, XP000872498 abstract	1-7
x	HOSOKAWA, M. ET AL: "Troglitazone inhibits bicarbonate secretion in rat and human duodenum" J. PHARMACOL. EXP. THER. (1999), 290(3), 1080-1084, XP008002057 page 1084, column 1, paragraph 3	1,2,4-6, 9,10
X	EP 0 861 666 A (TAKEDA CHEMICAL INDUSTRIES LTD) 2 September 1998 (1998-09-02) claims	1,2,4-6, 8,10
X	US 5 965 584 A (IKEDA HITOSHI ET AL) 12 October 1999 (1999-10-12) column 2, paragraph 2 - paragraph 4 column 3, paragraph 3 column 12, paragraph 3	1,2,4-6, 8,10
X	EP 0 930 076 A (SANKYO CO) 21 July 1999 (1999-07-21) claims	1,2,4-6, 8,10

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

International application No. PCT/GB 00/05006

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-7 and 10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. X Claims Nos.: 1, 2, 4-7,10 because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
*
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark on Protest The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees:

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1998)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1, 2, 4-7,10

Present claims 1, 2, 4-7 and 10 relate to compounds which are actually not well-defined: The use of the definitions "insulin sensitiser, such as compound I", "beta blocker", "thiazolidinedione insulin sensitiser", "an agent used in the treatment of the cardiac conditions associated with diabetes mellitus", "ACE inhibitor", "diuretic" and "endothelin antagonist" in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. An attempt is made to define the compounds by reference to a result to be achieved. The lack of clarity is such as to render a meaningful complete search impossible. Consequently, the search has been restricted to those parts relating to the compounds individually structurally identified by name in the claims, with due regard to the therapeutic application mentioned in the claims.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

Information on patent family members

.atlonal Application No PCT/GB 00/05006

	atent document		Publication		Patent family	Publication
	in search report		date		member(s)	date
MO	0027434	A	18-05-2000	AU	1063400 A	29-05-2000
				BR -	9915214 A	18-12-2001
				CN	1332638 T	23-01-2002
		•		EP	1128845 Al	05-09-2001
				WO	0027434 A1	18-05-2000
				NO	20012300 A	03-07-2001
EP	0861666	A	02-09-1998	EP	1174135 A2	23-01-2002
				EP	0861666 A2	02-09-1998
				AU	723097 B2	17-08-2000
				AU	5603496 A	09-01-1997
				CA	2179584 A1	21-12-1996
	•			CN	1145783 A	26-03-1997
	•			CZ	9601811 A3	15-01-1997
				EP	0749751 A2	27-12-1996
				HU	9601698 A2	28-05-1997
			·	JP	3148973 B2	26-03-2001
				JP	9067271 A	11-03-1997
			•	JP	10167986 A	23-06-1998
				NO	962606 A	23-12-1996
			•	NO	20004345 A	23-12-1996
				SK	79496 A3	08-01-1997
				TW	438587 B	07-06-2001
				ÜS	5965584 A	12-10-1999
•		-		ÜŠ	6150383 A	21-11-2000
				ÜŠ	6169099 B1	02-01-2001
				ÜS	6133293 A	17-10-2000
•				ÜŠ	6166042 A	26-12-2000
				ÜS	6214848 B1	10-04-2001
				ÜŠ	6166043 A	26-12-2000
				US	6150384 A	21-11-2000
				ÜŠ	6172089 B1	09-01-2001
				ÜS	6172090 B1	09-01-2001
				US	6121295 A	19-09-2000
			•	US	6156773 A	05-12-2000
	â			ÜS	6174904 B1	16-01-2001
				US	6121294 A	19-09-2000
				US	6225326 B1	01-05-2001
				US	6080765 A	27-06-2000
				US	6133295 A	17-10-2000
				US	6103742 A	15-08-2000
				US	6169100 B1	02-01-2001
				US	6329404 B1	11-12-2001
				US	6303640 B1	16-10-2001
				US	6211205 B1	03-04-2001
				US	6288090 B1	11-09-2001
				US	6232330 B1	15-05-2001
				US	6218409 B1	17-04-2001
				US	6323225 B1	27-11-2001
				US	6251924 B1	
				US	6239153 B1	26-06-2001
				US	6211206 B1	29-05-2001
				US		03-04-2001
				US	6211207 B1	03-04-2001
					5952356 A	14-09-1999
				US	6277869 B1	21-08-2001
				US	6274605 B1	14-08-2001
				US	6271243 B1	07-08-2001

Form PCT/ISA/210 (patent family annex) (July 1992)

Information on patent family members

utional Application No
PCT/GB 00/05006

Patent document cited in search report	Publication date		Patent family member(s)	Publication date
US 5965584 A	12-10-1999	US	6150383 A	21-11-2000
U3 5905564 A	12-10-1999	US	6169099 B1	02-01-2001
		US	·6133293 A	17-10-2000
		US	6166042 A	26-12-2000
		US	6214848 B1	10-04-2001
		US	6166043 A	26-12-2000
		US	6150384 A	21-11-2000
		US	6172089 B1	09-01-2001
		US	6172090 B1	09-01-2001
		US	6121295 A	19-09-2000
		US	6156773 A	05-12-2000
		ÜS	6103742 A	15-08-2000
		ÜS	6169100 B1	02-01-2001
• •		US	6329404 B1	11-12-2001
		US	6303640 B1	16-10-2001
	•	US	6211205 B1	03-04-2001
		US	6288090 B1	11-09-2001
		US	6232330 B1	15-05-2001
		US	6218409 B1	17-04-2001
		US	6323225 B1	27-11-2001
		US -	6251924 B1	26-06-2001
		US	6239153 B1	29-05-2001
		US	6211206 B1	03-04-2001
		US	6211207 B1	03-04-2001
		US	6277869 B1	21-08-2001
		US	6274605 B1	14-08-2001
•				07-08-2001
		US	6271243 B1	
		AU	723097 B2	17-08-2000
		AU	5603496 A	09-01-1997
		CA	2179584 A1	21-12-1996
		CN	1145783 A	26-03-1997
		CZ	9601811 A3	- 15-01-1997
		EP	. 1174135 A2	23-01-2002
		EP	0749751 A2	27-12-1996
		EP	0861666 A2	02-09-1998
		HU	9601698 A2	28-05-1997
		JP	3148973 B2	26-03-2001
		JP	9067271 A	11-03-1997
			10167986 A	
		JP		23-06-1998
		NO	962606 A	23-12-1996
		NO	20004345 A	23-12-1996
		SK	79496 A3	08-01-1997
		TW	438587 B	07-06-2001
		US	6174904 B1	16-01-2001
		US	6121294 A	19-09-200 0
		US	6225326 B1	01-05-2001
		US	6080765 A	27-06-2000
		ÜŠ	6133295 A	17-10-2000
		US	5952356 A	14-09-1999
		US	2002002186 A1	03-01-2002
		U3		
ED 0020076	21 07 1000	.VII	714610 D2	06-01-2000
EP 0930076 A	21-07-1999	AU	714618 B2	
		AU	3459597 A	09-02-1998
		EP	0930076 A1	21-07-1999
	•	HU	9903166 A2	28-09-2000
				15 00 1000
		NO	990166 A	15-03 - 199 9
		NO US	990166 A 2002013308 A1	31-01-2002

Information on patent family members

PCT/GB 00/05006

Patent document cited in search report	Publication date		Patent family member(s)	Publication date
EP 0930076 A	*	CN	1230122 A	29-09-1999
		CZ	9900102 A3	16-06-1999
		EP	1175902 A1	30-01-2002
		WO	9802183 A1	22-01-1998
		JР	10081632 A	31-03-1998